

STEPS FOR STARTING ENTYVIO



Please see Important Safety Information on pages 8, 9, and 10.



Getting treatment with ENTYVIO works a little differently than other prescriptions you may have used.

With a prescription for ENTYVIO from your doctor, you've taken the first step toward treating your moderate to severe Crohn's disease or ulcerative colitis. Because ENTYVIO is a biologic, the process might be different than what you're used to.

Read on to understand the **4 important steps** that will make your treatment journey with ENTYVIO as seamless as possible. If you ever have questions about the process, feel free to call us at 1-844-ENTYVIO (1-844-368-9846).

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on pages <u>8</u>, <u>9</u>, and <u>10</u>.











GET COVERAGE AUTHORIZATION

First, your health insurance provider must approve your coverage for ENTYVIO. These 2 steps are typically handled by your doctor's office.



Step 1: Benefits Verification

Your doctor's office will work with your insurance company to determine if ENTYVIO can be covered under your benefit plan.



Step 2: Prior Authorization

When required, your doctor's office will submit a request to your health insurance company to get approval for you to start treatment with ENTYVIO.

Keep in mind that the authorization process may take several weeks. Your doctor's office will be in touch if they need additional information.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on pages <u>8</u>, <u>9</u>, and <u>10</u>.



SIGN UP FOR ENTYVIOCONNECT

EntyvioConnect is a patient support program designed to help you throughout your entire treatment journey at no cost. Sign up at ENTYVIO.com/connect or call 1-844-ENTYVIO (1-844-368-9846)



One-on-One Nurse Support

You'll be paired with a Nurse Educator who can answer questions about Entyvio resources. While nurses can't give personalized medical advice, they can provide guidance to help you make the most out of every checkup you have with your doctor.



Financial and Insurance Help

A specialist can help you navigate co-pay assistance and outof-pocket costs. With our Co-Pay program, you may pay as little as \$5 per dose every 8 weeks.*





Ongoing Support

You can sign up to receive additional information or schedule helpful medication reminders. Whether you receive ENTYVIO at an infusion center, your doctor's office, or at home, our text reminders can help you stay on track.





Find an infusion center near you at **locator.infusioncenter.org**







PREPARE FOR YOUR FIRST INFUSION

Once your prescription is approved, it's time to get ready for your first treatment.



ENTYVIO is administered by a healthcare provider via an intravenous (IV) infusion (a needle inserted into a vein in your arm). Your doctor's office will help you choose the time and location for your first IV infusion. Depending on your coverage, you may receive an infusion at your doctor's office, an infusion center, a hospital, or your home.



Your doctor will monitor you during and after the infusion for side effects to see if you have a reaction to the treatment.

Get to your appointment early, and bring your insurance information and co-pay.

Safety Considerations

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on pages <u>8</u>, <u>9</u>, and <u>10</u>.



KNOW YOUR INFUSION SCHEDULE



Starting doses

You will receive ENTYVIO by infusion at Weeks 0, 2, and 6.



Maintenance doses

Starting at Week 14, you'll get 1 infusion every 8 weeks.

- In the first year, you'll receive 8 doses of ENTYVIO
- After the first year, you'll receive 6 doses of ENTYVIO

The infusion takes about **30 minutes**. Bring a good book, your favorite tunes, or something else to pass the time!

Uses of ENTYVIO® (vedolizumab)

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis
- moderately to severely active Crohn's disease







Important Safety Information for ENTYVIO® (vedolizumab)

- Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients.
- · ENTYVIO may cause serious side effects, including:
- Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. You may need treatment if you have an allergic reaction. Tell your healthcare provider or get immediate medical help if you get any of these symptoms during or after an infusion of ENTYVIO: rash, itching, swelling of your lips, tongue, throat or face, shortness of breath or trouble breathing, wheezing, dizziness, feeling hot, or palpitations (feel like your heart is racing).
- ENTYVIO may increase your risk of getting a serious infection. Before
 receiving and during treatment with ENTYVIO, tell your healthcare
 provider if you think you have an infection or symptoms of an infection,
 such as fever, chills, muscle aches, cough, shortness of breath, runny
 nose, sore throat, red or painful skin or sores on your body, tiredness,
 or pain during urination.

Important Safety Information for ENTYVIO (cont.)

- People with weakened immune systems can get progressive multifocal leukoencephalopathy (PML) (a rare, serious brain infection caused by a virus). Although unlikely while receiving ENTYVIO, a risk of PML cannot be ruled out. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML. Tell your healthcare provider right away if you have any of the following symptoms: confusion or problems thinking, loss of balance, change in the way you walk or talk, decreased strength or weakness on one side of the body, blurred vision, or loss of vision.
- Liver problems can happen in people who receive ENTYVIO. Tell your healthcare provider right away if you have any of the following symptoms: tiredness, loss of appetite, pain on the right side of your abdomen, dark urine, or yellowing of the skin and eyes (jaundice).
- The most common side effects of ENTYVIO include common cold, headache, joint pain, nausea, fever, infections of the nose and throat, tiredness, cough, bronchitis, flu, back pain, rash, itching, sinus infection, throat pain, and pain in extremities. These are not all the possible side effects of ENTYVIO. Call your healthcare provider for medical advice about side effects.

8



Important Safety Information for ENTYVIO (cont.)

 Before receiving ENTYVIO, tell your healthcare provider about all of your medical conditions, including if you: have or think you may have an infection or have infections that keep coming back; have liver problems; have tuberculosis (TB) or have been in close contact with someone with TB; have recently received or are scheduled to receive a vaccine; or if you are pregnant, breastfeeding, plan to become pregnant, or plan to breastfeed

Please see full <u>Prescribing Information</u>, including the <u>Medication Guide</u>, for ENTYVIO and talk with your healthcare provider.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Uses of ENTYVIO® (vedolizumab)

ENTYVIO is a prescription medicine used in adults for the treatment of:

- · moderately to severely active ulcerative colitis
- · moderately to severely active Crohn's disease

10







©2022 Takeda Pharmaceuticals U.S.A., Inc. 95 Hayden Ave., Lexington, MA 02421. 1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. TAKEDA and the TAKEDA logo are trademarks or registered trademarks of Takeda Pharmaceutical Company Limited. ENTYVIO is a trademark of Millennium Pharmaceuticals, Inc., registered with the U.S. Patent and Trademark Office and is used under license by Takeda Pharmaceuticals U.S.A., Inc.

US-VED-1815v1.0 12/22