



The only
gut-focused treatment*
for Crohn's and UC
that works right where U need it



*ENTYVIO helps block certain inflammation-causing cells from entering the gut.

Ask your healthcare
provider about ENTYVIO.



What is ENTYVIO?

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis (UC)
- moderately to severely active Crohn's disease (CD)

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).



➤ Focus U Want

Helps block inflammation right where it occurs: **In your gut**

*ENTYVIO helps block certain inflammation-causing cells from entering the gut.

[Go to page 3 to learn how ENTYVIO works](#)

➤ Results U Need

Lasting relief and **steroid-free remission** at 1 year[†]

Some people experienced **rapid symptom relief** as early as 6 weeks[†]

Individual results may vary.

[Go to page 4 for results with ulcerative colitis](#)

[Go to page 6 for results with Crohn's disease](#)

[†]Many patients taking ENTYVIO IV achieved remission, some without steroids, at Week 6 and Week 52. Remission is a measure to determine when a patient with Crohn's or UC is experiencing few to no symptoms and an improvement in their condition.

➤ Experience U Expect

More than 10 years of experience[‡]

#1 prescribed biologic for Crohn's and UC combined[§]

2 options for therapy:
ENTYVIO IV and the **ENTYVIO Pen**^{||}

[‡]ENTYVIO IV was FDA approved in May 2014.

[§]Quarterly analysis of claims data from January 2023-December 2025.

^{||}The ENTYVIO Pen is an option after at least 2 IV starter doses.

[Go to page 8 for administration information](#)

➤ Support U Deserve

Eligible ENTYVIO patients may pay as little as **\$0 per dose**

Growing coverage for the **ENTYVIO Pen**[†]

1-on-1 support through *EntyvioConnect* for people prescribed ENTYVIO

[Go to page 10 to see real patient experiences](#)

[Go to page 11 for support with *EntyvioConnect*](#)

[†]Coverage of the ENTYVIO Pen as of January 2025.

Speaking With Your Healthcare Provider

[Go to page 12 for a discussion guide](#)


SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).

**Focus U Want****How ENTYVIO works****How Crohn's and UC affect your gut****Inflammation**

Crohn's and UC are both immune system disorders that can cause inflammation in your gut.


**White blood cells**

White blood cells are part of your immune system. They help fight off infection and other diseases.

In people with Crohn's or UC, overactive white blood cells attack the healthy cells in your gut, causing inflammation.

ENTYVIO works right where U need it**Block**

ENTYVIO helps block certain inflammation-causing white blood cells from entering the gut.

**Reduce**

By helping to block white blood cells from entering the gut, ENTYVIO helps reduce inflammation caused by Crohn's and UC.

**Ask your healthcare provider if ENTYVIO is right for you**

Individual results may vary.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO


Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on page 14.

Ulcerative colitis


Rapid symptom relief at 6 weeks*

ENTYVIO helped **THREE TIMES** as many people achieve remission vs placebo



*Remission is a measure to determine when a patient with UC is experiencing a few to no symptoms and an improvement in their condition.

ENTYVIO helped many people with UC reach remission* at 1 year vs placebo



In clinical trials, ENTYVIO delivered results for adults with moderate to severe ulcerative colitis when other medicines had not worked well enough or could not be tolerated.

- The first study evaluated response to ENTYVIO IV versus placebo at Week 6. The second study included adults who had a previous response to ENTYVIO IV at Week 6 and evaluated remission at Week 52
- For the ENTYVIO Pen, a separate clinical study evaluated remission at Week 52 in adult patients with moderate to severe UC who were administered ENTYVIO subcutaneously vs placebo after at least 2 IV doses
- In the ENTYVIO clinical studies, clinical response and remission were defined and measured by improvement in the Mayo score, an assessment tool that measures stool frequency, rectal bleeding, appearance of the mucosal lining of the colon, and physician assessment

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).


Ulcerative colitis

Even more results U need



Nearly twice as many people achieved overall improvement


in their UC at 6 weeks with ENTYVIO vs placebo*



*Response is a measure to determine improvement of the patient's UC symptoms and conditions.

Steroid-free remission at 1 year


ENTYVIO IV helped more than twice as many people achieve remission without the use of steroids† at 1 year vs placebo



†Some patients were on steroids at the beginning of the study. ENTYVIO helped some of these patients achieve remission without the use of steroids at Week 52.

Superior to HUMIRA®

In a separate head-to-head study, ENTYVIO IV helped significantly more people achieve remission at 1 year vs HUMIRA‡



‡HUMIRA is a registered trademark of AbbVie Inc., North Chicago, IL. For more information related to adalimumab, please see AbbVie.com.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).

Crohn's disease




In clinical trials, ENTYVIO delivered results for adults with moderate to severe Crohn's disease when other medicines had not worked well enough or could not be tolerated.

- The first and second studies evaluated the number of patients who achieved remission on ENTYVIO versus placebo at Week 6. The third study enrolled adults who had a previous response to ENTYVIO at Week 6 and evaluated remission at Week 52
- For the ENTYVIO Pen, a separate clinical study evaluated remission at Week 52 in adult patients with moderate to severe Crohn's who were administered ENTYVIO subcutaneously vs placebo after at least 2 IV doses
- In the ENTYVIO clinical studies, clinical remission was defined and measured by improvement in the CDAI score, a tool that assesses the number of liquid or very soft stools, abdominal pain, and other measures of disease activity

Rapid symptom relief at 6 weeks*

ENTYVIO helped more than twice as many people achieve remission at 6 weeks vs placebo



*Remission is a measure to determine when a patient with Crohn's is experiencing a few to no symptoms and an improvement in their condition.

Ask your healthcare provider if ENTYVIO is right for you

Individual results may vary.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.


Please see Important Safety Information on [page 14](#).

Crohn's disease

Even more results U need




ENTYVIO helped many people with Crohn's **reach remission*** at 1 year vs placebo



*Remission is a measure to determine when a patient with Crohn's is experiencing a few to no symptoms and an improvement in their condition.

Steroid-free remission at 1 year

ENTYVIO IV helped twice as many people achieve remission without depending on steroids[†] at 1 year vs placebo



[†]Some patients were on steroids at the beginning of the study. ENTYVIO helped some of these patients achieve remission without the use of steroids at Week 52.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).

Experience U Expect

Administration options

2 options for maintenance therapy

You and your healthcare provider can choose between ENTYVIO IV and the ENTYVIO Pen (available after at least 2 IV starter doses).



ENTYVIO intravenous (IV) treatment

ENTYVIO IV treatment is administered in an office, infusion center, or at home with medical support.

- A healthcare professional administers ENTYVIO by IV infusion and stays to monitor your treatments
- The ENTYVIO IV is delivered by a needle in the arm and takes about 30 minutes
- After your starting doses, the infusion is given every 8 weeks



The ENTYVIO Pen

The ENTYVIO Pen provides the same medicine in a form that may better fit your lifestyle.

- The ENTYVIO Pen can be administered in several areas of the body
- You or your caregiver inject ENTYVIO under the skin every 2 weeks after at least 2 IV starter doses
- Your healthcare provider will help train you on how to use this single-dose, prefilled pen. You can also receive virtual or in-home injection training before you start
- After you are prescribed the ENTYVIO Pen, a specialty pharmacy will reach out to you regarding coordination and delivery of your prescription

Feel confident in your treatment choice—no matter the administration option

Whichever option you and your healthcare provider choose, you can feel confident knowing the ENTYVIO in your blood will consistently stay above a certain level—and that switching to the ENTYVIO Pen does not lead to lower ENTYVIO drug amounts in your body.*

The ENTYVIO Pen and ENTYVIO IV showed similar overall safety profiles, with the exception of injection site reactions, which were reported in 10% (11/106) of patients using the ENTYVIO Pen.

*In the study of UC patients, the blood level of ENTYVIO was an exploratory data point and cannot be used to determine efficacy or safety.

Use our infusion center finder to find a location that's convenient for you.
Go to **infusionfinder.ENTYVIOHCP.com**.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on page 14.

Intravenous infusion dosing schedule

Starting treatment schedule



Week 0



Week 2



Week 6

You will receive ENTYVIO by infusion at **Weeks 0, 2, and 6.**

Self-injection dosing schedule

Starting treatment schedule



Week 0



Week 2

You will receive ENTYVIO by infusion at **Weeks 0 and 2.**

See page 8 to learn more about infusions and where you can receive yours.

Your healthcare provider will determine if it's appropriate to transition to self-administration with the ENTYVIO Pen after at least 2 IV doses.

Maintenance schedule

After your initial doses, you will continue to receive **ENTYVIO by IV every 8 weeks.**



Every 8 Weeks

- In the first year, you'll receive **8 doses of ENTYVIO**
- After the first year, you'll receive **6 doses each year**

Maintenance schedule



After your initial doses, you or a caregiver will use the ENTYVIO Pen to inject ENTYVIO in place of your next scheduled IV dose, and then **every 2 weeks** up to 26 times per year.

See the full Instructions for Use at [ENTYVIO.com/IFU](https://www.entyvio.com/IFU).

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).

Support U Deserve

What real patients are saying

Living with Crohn's or UC brings its challenges. But for these patients, ENTYVIO has made a difference. Individual results may vary.

"My doctor explained to me that ENTYVIO helps block a type of inflammation-causing white blood cell from entering the gut. I really liked that."



Shivani: Living with moderate to severe UC

"After I began IV treatment with ENTYVIO, my symptoms started to improve and I began to feel better. I hadn't even realized how much I had become accustomed to having pain in my gut."



Ben: Living with moderate to severe Crohn's

Entyvio Connections

Connect with a real ENTYVIO patient now

Sign up for the ENTYVIO Connections program to talk with one of our Patient Ambassadors about ENTYVIO and their experiences with Crohn's and UC.

Call **1-833-492-8847** or visit **[ENTYVIO.com/mentor](https://www.ENTYVIO.com/mentor)** to register for ENTYVIO Connections.



Donna: Living with moderate to severe Crohn's

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).

Support U Deserve

EntyvioCONNECT

Get ongoing support with *EntyvioConnect*

Living with Crohn's or UC can be difficult, but managing your treatment shouldn't be. The caring team of experts at *EntyvioConnect* is available to guide you with resources for both ENTYVIO IV and the ENTYVIO Pen.

Once you're prescribed ENTYVIO, you're eligible for ongoing support with *EntyvioConnect*, including treatment reminders, useful resources, healthy recipes, and much more.

One-on-one nurse support

We can pair you with a Nurse Educator who can answer your questions about ENTYVIO resources and help you make the most out of every conversation with a healthcare provider. They cannot, however, provide medical advice.

If you are prescribed the ENTYVIO Pen, virtual and in-person nurse injection training is available through *EntyvioConnect*.

Pay as little as \$0 per dose*

There's currently more than 90% commercial coverage for ENTYVIO IV and growing coverage for the ENTYVIO Pen. With our Co-Pay Program, you may pay as little as \$0 per dose.* An *EntyvioConnect* Case Manager can help you navigate the insurance and co-pay assistance process.

*Must meet eligibility requirements.

SELECTED SAFETY CONSIDERATIONS WITH ENTYVIO

Do not receive ENTYVIO if you have had an allergic reaction to ENTYVIO or any of its ingredients. Infusion-related and serious allergic reactions can happen while you are receiving ENTYVIO or several hours after treatment. ENTYVIO may increase your risk of getting a serious infection.

Please see Important Safety Information on [page 14](#).



The ENTYVIO Start Program

No cost.

Zero enrollment delays.

For up to 3 years.

With **the ENTYVIO Start Program**, it's easier than ever to start and stay on ENTYVIO with the customized, free support you deserve—available for both IV and the ENTYVIO Pen. Ask your healthcare provider or an *EntyvioConnect* Case Manager for details.

Eligibility requirements apply.

Terms and Conditions apply.

Get connected

Call 1-844-ENTYVIO (1-844-368-9846)

or sign up at [ENTYVIO.com/Connect](https://www.ENTYVIO.com/Connect).



Talking to your healthcare provider

The questions below can help get your conversation started as you and your healthcare provider explore potential treatment options for your condition. Fill out this page and bring it with you to your next appointment.

How many times have you had to refill your prescription of steroids in the last year?

- ☐ Never ☐ Once ☐ Monthly ☐ More

How often do you currently take steroids to control flare-ups, as prescribed by your healthcare provider?

- ☐ Daily ☐ Weekly ☐ Monthly
☐ Once every 2 months ☐ Once every 3 months

How many bowel movements do you have a day?

- ☐ 1 ☐ 2 ☐ 3 ☐ 4 ☐ 5
☐ 6 ☐ 7 ☐ 8 ☐ 9 ☐ 10+

Have you experienced any weight loss?

- ☐ Yes ☐ No

This is a sample dialogue. It is not, nor is it intended to be, a medical evaluation, examination, advice, consultation, diagnosis, or treatment. Always consult your healthcare professional for all medical and health-related matters.

How often do you experience any of the following symptoms?

Abdominal pain or cramping

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Diarrhea

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Rectal bleeding

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Accidents

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Constipation

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Straining during bowel movements

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Lack of energy

- ☐ Never ☐ Occasionally ☐ Sometimes
☐ Often ☐ Always

Please indicate how you feel using the following scale



Are your current treatments working well enough?

- ☐ Yes ☐ No

Your healthcare provider can help you find a treatment that works best for you.

Questions to ask your doctor

Here are some examples of questions you may want to ask your doctor as you work together to find the best treatment for your specific needs.

How is a biologic medication different from other medications?

How does ENTYVIO work to help fight inflammation?

How do you and I choose between IV infusion and self-administered injection?

How will I know if the treatment is working for me?

What goals should I set for my treatment?

Could ENTYVIO be right for me?

IMPORTANT SAFETY INFORMATION

- **Do not receive ENTYVIO® if you** have had an allergic reaction to ENTYVIO or any of its ingredients.
- **ENTYVIO may cause serious side effects, including:**
 - **Infusion-related and serious allergic reactions** can happen while you are receiving ENTYVIO or several hours after treatment. You may need treatment if you have an allergic reaction. Tell your healthcare provider or get immediate medical help if you get any of these symptoms during or after an infusion of ENTYVIO: rash, itching, swelling of your lips, tongue, throat or face, shortness of breath or trouble breathing, wheezing, dizziness, feeling hot, or palpitations (feel like your heart is racing).
 - **Infections.** ENTYVIO may increase your risk of getting a serious infection. Before and during treatment with ENTYVIO, tell your healthcare provider if you think you have an infection or symptoms of an infection, such as fever, chills, muscle aches, cough, shortness of breath, runny nose, sore throat, red or painful skin or sores on your body, tiredness, or pain during urination. If your healthcare provider feels that you are at risk for tuberculosis (TB), you may be treated with medicine for TB before you begin treatment with ENTYVIO.
 - **Progressive Multifocal Leukoencephalopathy (PML).** People with weakened immune systems can get progressive multifocal leukoencephalopathy (PML) (a rare, serious brain infection caused by a virus). Although unlikely while receiving ENTYVIO, a risk of PML cannot be ruled out. PML can result in death or severe disability. There is no known treatment, prevention, or cure for PML. Tell your healthcare provider right away if you have any of the following symptoms: confusion or problems thinking, loss of balance, change in the way you walk or talk, decreased strength or weakness on one side of the body, blurred vision, or loss of vision.
 - **Liver problems** can happen in people who receive ENTYVIO. Tell your healthcare provider right away if you have any of the following symptoms: tiredness, loss of appetite, pain on the right side of your abdomen, dark urine, or yellowing of the skin and eyes (jaundice).
- **The most common side effects of ENTYVIO include:** common cold, headache, joint pain, nausea, fever, infections of the nose and throat, tiredness, cough, bronchitis, flu, back pain, rash, itching, sinus infection, throat pain, pain in extremities, and with injections under the skin: pain, swelling, itching, hives, bruising, rash, or redness at the injection site. These are not all the possible side effects of ENTYVIO. Call your healthcare provider for medical advice about side effects.
- **Before receiving ENTYVIO, tell your healthcare provider about all of your medical conditions, including if you:** have or think you may have an infection or have infections that keep coming back; have liver problems; have TB or have been in close contact with someone with TB; have recently received or are scheduled to receive a vaccine; or if you are pregnant, breastfeeding, plan to become pregnant, or plan to breastfeed.
- **Tell your healthcare provider about all the medicines you take, especially if you take or have recently taken** Tysabri (natalizumab), Tyruko (natalizumab-sztn), a Tumor Necrosis Factor (TNF) blocker medicine, a medicine that weakens your immune system (immunosuppressant), or corticosteroid medicine.

Please see accompanying **full Prescribing Information**, including the **Medication Guide**, for ENTYVIO and talk with your healthcare provider.

ENTYVIO is available for:

- **intravenous (IV) infusion: 300 mg vedolizumab.**
- **subcutaneous (SC) injection: 108 mg vedolizumab.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

What is ENTYVIO (vedolizumab)?

ENTYVIO is a prescription medicine used in adults for the treatment of:

- moderately to severely active ulcerative colitis (UC).
- moderately to severely active Crohn's disease (CD).

It is not known if ENTYVIO is safe and effective in children under 18 years of age.



**Ask your healthcare
provider about getting the
results U want
with ENTYVIO**

Sign up for more information or call us at 1-844-ENTYVIO (1-844-368-9846)

*ENTYVIO helps block certain inflammation-causing cells from entering the gut.

Please see Important Safety Information on page 14 and [Prescribing Information](#).



©2026 Takeda Pharmaceuticals U.S.A., Inc., 500 Kendall Street, Cambridge, MA 02142. 1-877-TAKEDA-7 (1-877-825-3327). All rights reserved. TAKEDA and the TAKEDA logo are registered trademarks of Takeda Pharmaceutical Company Limited. ENTYVIO and the ENTYVIO logo are registered trademarks of Millennium Pharmaceuticals, Inc. All other trademarks are the property of their respective owners.

US-VED-3699v3.0 04/26